

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11960 - Goat Anti-PLA2G7 (aa53-64) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PLA2G7, phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma), LDL-PLA2, LP-PLA2, PAFAD, PAFAH, 1-alkyl-2-acetylglycerophosphocholine esterase, 2-acetyl-1-alkylglycerophosphocholine esterase, LDL-PLA(2), LDL-associated phospholipase A2, PAF 2-acylhydrolase, PAF

acetylhydrolase, gVIIA-PLA2, group-VIIA phospholipase A2, lipoprotein-associated phospholipase A2, platelet-activating factor acetylhydrolase **Official Symbol:** PLA2G7

Accession Number(s): NP_005075.3

Human GeneID(s): 7941

Important Comments: Reported variants represent identical protein: NP_005075.3, NP_001161829.1

Immunogen

Peptide with sequence C-QTKIPRGNGPYS, from the internal region of the protein sequence according to NP_005075.3.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Preliminary experiments gave an approx 100kDa band in Human Liver lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 50.1kDa according to NP_005075.3. The 100kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Does the observed molecular weight correspond to a glycosylated form?

Species Reactivity

Tested: Expected from sequence similarity: Human