

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06969 - Goat Anti-PLK1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** PLK1, polo-like kinase 1 (Drosophila), HGNC:9077, PLK, STPK13, cell cycle regulated protein kinase polo (Drosophia)-like kinase, polo like kinase, polo-like

kinase, polo-like kinase (Drosophila)

Official Symbol: PLK1

Accession Number(s): NP\_005021.2

Human GeneID(s): 5347

Non-Human GenelD(s): 18817 (mouse), 25515 (rat)

### **Immunogen**

Peptide with sequence C-DKLLSSRSASNRLK, from the C Terminus of the protein sequence according to NP\_005021.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 65kDa band observed in lysates of cell line K562 (calculated MW of 68.3kDa according to NP\_005021.2). Recommended concentration: 1-3µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB06969 (1 $\mu$ g/ml) staining of K562 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.