

## **International Office**

#### Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08403 - Goat Anti-PNPLA2 / Desnutrin Antibody

Size: 100µg specific antibody in 200µl

# **Target Protein**

Principal Names: PNPLA2, desnutrin, patatin-like phospholipase domain containing 2, 1110001C14Rik, ATGL, DKFZp667M109, FP17548, PEDF-R, TTS-2.2, TTS2, adipose triglyceride lipase, pigment epithelium-derived factor, transport-secretion protein 2.2, triglyceride hydrolase Official Symbol: PNPLA2 Accession Number(s): NP\_065109.1 Human GenelD(s): 57104

#### Immunogen

Peptide with sequence C-HFNSKDELIQAN, from the internal region of the protein sequence according to NP\_065109.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 60kDa band observed in Human Heart lysates (calculated MW of 55.3kDa according to NP\_065109.1). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour. This antibody has been successfully used in WB on Mouse, PMID: 29175050.

**Immunofluorescence:** This antibody has been successfully used in IF on Human in the following papers, PMID: 33160079.

## **Species Reactivity**

Tested: Human, Mouse Expected from sequence similarity: Human, Mouse, Rat, Cow

## **Specific References**

This antibody has been successfully used in IF on Human:

Anne Gemmink, Sabine Daemen, Bram Brouwers, Joris Hoeks, Gert Schaart, Kèvin Knoops, Patrick Schrauwen, Matthijs K C Hesselink

Decoration of myocellular lipid droplets with perilipins as a marker for in vivo lipid droplet dynamics: A super-resolution microscopy study in trained athletes and insulin resistant individuals

Biochim Biophys Acta Mol Cell Biol Lipids. 2020 Nov 4;1866(2):158852. PMID: 33160079

## This antibody has been successfully used in WB on Mouse:

Ramseyer VD, Kimler VA, Granneman JG. Vacuolar protein sorting 13C is a novel lipid droplet protein that inhibits lipolysis in brownadipocytes. Mol Metab. 2018 Jan;7:57-70. PMID: 29175050

New Contraction	250kDa 150kDa 100kDa
2	75kDa
	50kDa
	37kDa
	25kDa
Via.	20kDa
14	15kDa

EB08403 (0.03µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.