



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05034 - Goat Anti-PPP1R15A / GADD34 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PPP1R15A, GADD34, protein phosphatase 1, regulatory (inhibitor) subunit 15A, growth arrest and DNA-damage-inducible 34, protein phosphatase 1, regulatory subunit 15A

Official Symbol: PPP1R15A

Accession Number(s): NP_055145.3

Human GeneID(s): [23645](#)

Immunogen

Peptide with sequence C-AAALDLSGRRG, from the C Terminus of the protein sequence according to NP_055145.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 75kDa band observed in lysates of cell line HepG2 (calculated MW of 73.4kDa according to NP_055145.2). Recommended concentration: 0.3-1µg/ml.

Primary incubation 1 hour at room temperature. **Negative Control:** KLY (Kelly Human neuroblastoma) cell lysate.

IHC: In paraffin embedded Human Liver shows cytoplasmic staining consistent with ER in the hepatocytes. Recommended concentration: 2-4µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nuclei of HeLa and HepG2 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HepG2 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human




EB05034 (0.3 μ g/ml) staining of HepG2 (A) and negative control KLY (B) lysate (35 μ g protein in RIPA buffer) Detected by chemiluminescence.




EB05034 (2 μ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.



EB05034 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing strong nuclear and some cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB05034 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing strong nuclear and some cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB05034 Flow cytometric analysis of paraformaldehyde fixed HepG2 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (0.4ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.