



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09395 - Goat Anti-PPP4C Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PPP4C, protein phosphatase 4 (formerly X), catalytic subunit, PP4, PPH3, PPX, Protein phosphatase 4, catalytic subunit

**Official Symbol:** PPP4C

**Accession Number(s):** NP\_002711.1

**Human GeneID(s):** [5531](#)

**Non-Human GeneID(s):** 171366 (rat)

### Immunogen

Peptide with sequence C-PQETRGIPSKKP, from the C Terminus of the protein sequence according to NP\_002711.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.


**Western blot:** Approx 35kDa band observed in HEK293, A431 and HeLa lysates and in Rat Spleen and Lung lysates (calculated MW of 35.1kDa according to NP\_002711.1).  
Recommended concentration: 0.1-0.3µg/ml.

**IHC:** In paraffin embedded Human Prostate shows nuclear staining of secretory cells in the gland. Recommended concentration, 2-4µg/ml.


### Species Reactivity

**Tested:** Human, Rat


**Expected from sequence similarity:** Human, Rat, Dog



EB09395 (0.3 $\mu$ g/ml) staining of HEK293 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.



EB09395 (0.3 $\mu$ g/ml) staining of Rat Spleen (A) and Lung (B) lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB09395 (2.5µg/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.