

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10761 - Goat Anti-PRDM14 (aa414-427) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** PRDM14, PR domain containing 14, MGC59730, PFM11, PR domain zinc finger protein 14, PR domain-containing protein 14, PR-domain zinc finger protein 14

Official Symbol: PRDM14

Accession Number(s): NP\_078780.1

Human GeneID(s): 63978

Non-Human GenelD(s): 383491 (mouse), 100365184 (rat)

#### **Immunogen**

Peptide with sequence YRDKHLKYTPCVDK, from the internal region of the protein sequence according to NP\_078780.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Preliminary experiments gave an approx 40kDa band in Human Heart lysates after 0.1-1μg/ml antibody staining and this band was successfully blocked by incubation with the immunizing peptide. Preliminary experiments also gave an approxand an approx 48-50kDa band in Human Lung and Kidney lysates and in nuclear lysates of cell liines Jurkat, HEK293 and MCF7. Please note that currently we cannot find an explanation in the literature for these bands we observe given the calculated size of 64.1kDa according to Human NP\_078780.1.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

#### **Species Reactivity**


Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB10761 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control:

Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB10761 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.