

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB13058 - Goat Anti-prepro-transforming growth factor beta-2 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** TGFB2, transforming growth factor beta 2, LDS4, TGF-beta2, BSC-1 cell growth inhibitor, G-TSF, cetermin, glioblastoma-derived T-cell suppressor factor;, polyergin, prepro-transforming growth factor beta-2, transforming growth factor, beta 2

Official Symbol: TGFB2

Accession Number(s): NP\_001129071.1; NP\_003229.1

Human GeneID(s): 7042

Important Comments: This antibody is expected to recognize the part representing the

latency-associated peptide.

#### **Immunogen**

Peptide with sequence C-STYTSGDQKTIKSTR, from the internal region of the protein sequence according to NP\_001129071.1; NP\_003229.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 55kDa band observed in Human Breast cancer lysates (calculated MW of 50.6kDa according to NP\_001129071.1;). This molecular weight is routinely observed by other sources. Recommended concentration: 0.1-0.3µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB13058 (0.3 $\mu$ g/ml) staining of Human Breast cancer lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.