

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11563 - Goat Anti-PRMT7 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** [Myelin basic protein]-arginine N-methyltransferase PRMT7, FLJ10640, histone-arginine N-methyltransferase PRMT7, KIAA1933, myelin basic protein-arginine

N-methyltransferase, OTTHUMP00000174863, PRMT7, protein arginine

methyltransferase 7, protein arginine N-methyltransferase 7

Official Symbol: PRMT7

Accession Number(s): NP\_061896.1; NP\_001171753.1

Human GeneID(s): 54496

Non-Human GenelD(s): 214572 (mouse), 361402 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_061896.1; NP\_001171753.1).

## **Immunogen**

Peptide with sequence C-PRFGEINDQDRTDR, from the internal region of the protein sequence according to NP\_061896.1; NP\_001171753.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 75kDa band observed in lysates of cell line HeLa and of Mouse Spleen (calculated MW of 78.5kDa according to Human NP\_061896.1 and 78.3kDa according to Mouse NP\_663379.1). Recommended concentration: 0.3-1µg/ml.

## **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa

EB11563 (0.3 $\mu$ g/ml) staining of HeLa lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.