



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11324 - Goat Anti-Progranulin aa248-259 / Granulin-3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** acrogranin, GEP, GP88, granulin, granulin-epithelin, granulins, PC cell-derived growth factor, PCDGF, PEPI, PGRN, proepithelin, progranulin, GRN

**Official Symbol:** GRN

**Accession Number(s):** NP\_002078.1

**Human GeneID(s):** [2896](#)

**Important Comments:** This antibody is expected to recognize the C-Terminal part of Granulin-3.

### Immunogen

Peptide with sequence PQDTVCDLIQSK, from the internal of the protein sequence according to NP\_002078.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 75kDa band observed in lysates of cell lines A431, A549 and HeLa (calculated MW of 63.5kDa according to NP\_002078.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (He and Bateman, Cancer Res. 1999 Jul 1;59(13):3222-9. PMID: 10397269). Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB11324 (1µg/ml) staining of A431 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.