

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09682 - Goat Anti-PROKR2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PROKR2, prokineticin receptor 2, GPR73L1, GPR73b, GPRg2, KAL3, PKR2, dJ680N4.3, G protein-coupled receptor 73-like 1, OTTHUMP00000030192

Official Symbol: PROKR2

Accession Number(s): NP_658986.1

Human GenelD(s): 128674

Non-Human GenelD(s): 192649 (rat)

Immunogen

Peptide with sequence C-NGNTSFTPNFN, from the internal region of the protein sequence according to NP_658986.1.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 45kDa band observed in Human Brain (Cerebellum), Testis and Thyroid Gland and in Rat Brain, Stomach and Testis lysates (calculated MW of 44.0kDa according to Human NP_658986.1 and 47.9kDa according to Rat NP_620434.1). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Brain (Cortex). Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human, Rat


Expected from sequence similarity: Human, Rat



EB09682 (0.1μg/ml) staining of Human Cerebellum (A), Rat Brain (B) and Rat Testis (C) lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB09682 (1 μ g/ml) staining of Rat Stomach lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB09682 (3.75 μ g/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.