



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11821 - Goat Anti-PROM1 / CD133 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AC133, antigen AC133, CD133, CORD12, hematopoietic stem cell antigen, hProminin, MCDR2, OTTHUMP00000217744, OTTHUMP00000217745, OTTHUMP00000217746, PROM1, prominin 1, prominin-1, prominin-like 1, prominin-like protein 1, PROML1, RP41, STGD4

Official Symbol: PROM1

Accession Number(s): NP_006008.1; NP_001139319.1; NP_001139324.1; NP_001139323.1; NP_001139322.1; NP_001139321.1

Human GeneID(s): [8842](#)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_006008.1; NP_001139319.1; NP_001139324.1; NP_001139323.1; NP_001139322.1; NP_001139321.1). Reported variants represent identical protein: NP_001139320.1, NP_001139319.1

Immunogen

Peptide with sequence RHATPTTRGCVSN, from the internal region of the protein sequence according to NP_006008.1; NP_001139319.1; NP_001139324.1; NP_001139323.1; NP_001139322.1; NP_001139321.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB11821 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).