

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09299 - Goat Anti-PSPH Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** PSPH, phosphoserine phosphatase, PSP, L-3-phosphoserine phosphatase, O-phosphoserine phosphohydrolase, OTTHUMP00000025059, PSPase

Official Symbol: PSPH

Accession Number(s): NP\_004568.2

Human GeneID(s): 5723

Non-Human GenelD(s): 100678 (mouse), 304429 (rat)

## **Immunogen**

Peptide with sequence C-RQQVKDNAKWYITD, from the C Terminus of the protein sequence according to NP\_004568.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 26kDa band observed in Human Brain (Hippocampus) and in Mouse brain lysates (calculated MW of 25.0kDa according to Human NP\_004568.2 and 25.1kDa according to Mouse NP\_598661.1). Recommended concentration: 0.05-1µg/ml.

## **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat,



EB09299 (0.05μg/ml) staining of Mouse Brain lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.