



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05314 - Goat Anti-RAC2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: RAC2, ras-related C3 botulinum toxin substrate 2 (rho family, small GTP binding protein Rac2), EN-7, Ras-related C3 botulinum toxin substrate 2, rho family, small GTP binding protein Rac2, Ras-related C3 botulinum toxin substrate 3 (rho family, small GTP-binding protein Rac2), Gx, HSPC022, small G protein

Official Symbol: RAC2

Accession Number(s): NP_002863.1

Human GeneID(s): [5880](#)

Immunogen

Peptide with sequence PGPTRQQKRACSL, from the C Terminus of the protein sequence according to NP_002863.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 22kDa band observed in lysates of cell line Molt4 (calculated MW of 21.4kDa according to NP_002863.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation 1 hour at room temperature.

IHC: In paraffin embedded Human Tonsil shows staining in mantle zone and some in the interfollicular area. Unstained in germinal centre. Recommended concentration, 2-4µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB05314 (0.1µg/ml) staining of Molt4 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB05314 (2µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining.