

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09551 - Goat Anti-RAP80 Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: UIMC1, ubiquitin interaction motif containing 1, RAP80, X2HRIP110, OTTHUMP00000161441, receptor associated protein 80, retinoid x receptor interacting protein

Official Symbol: UIMC1 Accession Number(s): NP_057374.3 Human GenelD(s): <u>51720</u> Non-Human GenelD(s): 20184 (mouse), 290997 (rat)

Immunogen

Peptide with sequence C-TRRQKEAKTKSDSG, from the internal region of the protein sequence according to NP_057374.3.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments gave an approx 60kDa band in Human Heart and Spleen lysates and in lysates of cell lines A431, HeLa and HepG2 after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 79.7kDa according to NP_057374.3. The 60kDa band was successfully blocked by incubation with the immunizing peptide. **IHC:** Paraffin embedded Human Testis. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

Auto-generated 12/05/2024





EB09551 (3.75 μ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.