

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06672 - Goat Anti-RARA Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: RARA, RAR, NR1B1, retinoic acid receptor, alpha, Retinoic acid receptor, alpha polypeptide, nucleophosmin-retinoic acid receptor alpha fusion protein NPM-RAR long form, nucleophosmin-retinoic acid receptor alpha fusion protein NPM-RAR short form

Official Symbol: RARA

Accession Number(s): NP_000955.1; NP_001138773.1; NP_001019980.1; NP_001138774.1

Human GeneID(s): [5914](#)

Important Comments: This antibody is expected to recognize all three reported isoforms (NP_000955.1; NP_001019980.1; NP_001138774.1). Please note the epitope is from the mouse/rat sequence. Human sequence is SPSLSPSSNRSSPATHSP and thus highly similar. Reported variants NP_000955.1 and NP_001138773.1 represent identical proteins.

Immunogen

Peptide with sequence C-SPSLSPSSHRSSPATQSP, from the C Terminus of the protein sequence according to NP_000955.1; NP_001138773.1; NP_001019980.1; NP_001138774.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 55kDa band observed in Human Breast and Breast cancer lysates (calculated MW of 50.8kDa according to NP_000955.1 and NP_001019980.1).
Recommended concentration: 0.5-2µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB06672 (1µg/ml) staining of Human Breast (A) and Breast cancer (B) lysate (35µg protein in RIPA buffer).
Primary incubation was 1 hour. Detected by chemiluminescence.