

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05421 - Goat Anti-RGS1 / 1R20 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** RGS1, 1R20, regulator of G-protein signalling 1, BL34, IER1, IR20, 1R20, Regulator of G-protein signaling 1, B-cell activation protein BL34, early response protein 1R20, immediate-early response 1, B-cell specific

**Official Symbol:** RGS1

**Accession Number(s):** NP\_002913.3

**Human GeneID(s):** [5996](#)

**Non-Human GeneID(s):** 50778 (mouse)

### Immunogen

Peptide with sequence C-NLLNDLQANSLK, from the C Terminus of the protein sequence according to NP\_002913.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 25kDa band seen in HepG2 lysate. Recommended for use at 0.5-2µg/ml.

**IHC:** Paraffin embedded Human Thymus and Small Intestine. Recommended concentration: 2.5µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB05421 staining (1µg/ml) of HepG2 lysate (RIPA buffer, 30µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



EB05421 (2.5µg/ml) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05421 (2.5µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.