

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06732 - Goat Anti-RICTOR Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** Rictor, KIAA1999, AVO3, MGC39830, likely ortholog of mouse TORC2-specific protein AVO3 (S. cerevisiae), rapamycin-insensitive companion of mTOR, mAVO3, pianissimo, DKFZp686B11164, RICTOR, RPTOR independent companion of MTOR, complex 2, PIA, mAVO3, TORC2-specific protein AVO3

Official Symbol: RICTOR

Accession Number(s): NP\_689969.2

Human GenelD(s): 253260

Non-Human GenelD(s): 78757 (mouse)

### Immunogen

Peptide with sequence C-KQPIVDTSAES, from the C Terminus of the protein sequence according to NP\_689969.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.


Western blot: Preliminary experiments gave an approx 38kDa band in Human Lung Carcinoma A549 cell line and cervix epitheloid carcinoma cell line HeLa lysates after 0.1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 192kDa according to NP\_689969.2. The 38kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**IHC:** In paraffin embedded Kidney shows strong cytoplasmic staining of epithelial cells in distal convoluted tubules. Recommended concentration, 2-3µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Dog



EB06732 (2.5 $\mu$ g/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.