

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07058 - Goat Anti-ROBO1 / DUTT1 (Internal) Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** ROBO1, DUTT1, roundabout, axon guidance receptor, homolog 1 (Drosophila), HGNC:10249, FLJ21882, SAX3, axon guidance receptor, roundabout 1,

MGC131599, MGC133277
Official Symbol: ROBO1

Accession Number(s): NP\_002932.1; NP\_598334.1

Human GenelD(s): 6091

Non-Human GenelD(s): 19876 (mouse), 58946 (rat)

Important Comments: Not expected to cross-react with ROBO2 or ROBO3.

## **Immunogen**

Peptide with sequence C-GDVDLSNKINEMK, from the internal region of the protein sequence according to NP\_002932.1; NP\_598334.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

IHC: Paraffin embedded Human Brain (Cortex) and Kidney. Recommended


concentration: 5-6µg/ml.

Immunofluorescence: Strong expression of the protein seen in the Plasma membrane and Golgi/ER of HeLa and U2OS cells. Recommended concentration: 10µg/ml.

# **Species Reactivity**


Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB07058 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton.


Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Plasma membrane and Golgi staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




EB07058 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Plasma membrane and ER staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




EB07058 (5µg/ml) staining of paraffin embedded Human Cortex. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07058 Negative Control showing staining of paraffin embedded Human Cortex, with no primary antibody.



EB07058 (6μg/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07058 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.