

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08028 - Goat Anti-RPL17 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: RPL17, ribosomal protein L17, MGC117162, rpL23, 60S ribosomal

protein L17, gene encoding putative NFkB activating protein

Official Symbol: RPL17

Accession Number(s): NP\_000976.1; NP\_001030178.1

Human GeneID(s): 6139

Non-Human GenelD(s): 319195 (mouse), 291434 (rat)

Important Comments: Both reported variants represent identical protein (NP\_000976.1;

NP\_001030178.1)

# Immunogen

Peptide with sequence RYSLDPENPTKSC, from the N Terminus of the protein sequence according to NP\_000976.1; NP\_001030178.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 23kDa band observed in Human Prostate lysates (calculated MW of 21.4kDa according to NP\_000976.1). Recommended concentration: 0.3-1µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

### **Specific Reference**

Smolock EM, Korshunov VA, Glazko G, Qiu X, Gerloff J, Berk BC.

Ribosomal protein L17, RpL17, is an inhibitor of vascular smooth muscle growth and carotid intima formation.

Circulation. 2012 Nov 13;126(20):2418-27.

PMID: 23065385

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB08028 (0.3 $\mu$ g/ml) staining of Human Prostate lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.