



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11513 - Goat Anti-S100A8 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** 60B8AG, CAGA, calgranulin A, calgranulin-A, calprotectin L1L subunit, CFAG, CGLA, CP-10, cystic fibrosis antigen, L1Ag, leukocyte L1 complex light chain, MA387, MIF, migration inhibitory factor-related protein 8, MRP8, MRP-8, NIF, OTTHUMP00000015329, P8, protein S100-A8, S100 calcium binding protein A8, S100 calcium-binding protein A8 (calgranulin A), S100A8, urinary stone protein band A

**Official Symbol:** S100A8

**Accession Number(s):** NP\_002955.2

**Human GeneID(s):** [6279](#)

### Immunogen

Peptide with sequence C-KGADVWFKELDINT, from the internal region of the protein sequence according to NP\_002955.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**IHC:** Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the nuclei and cytoplasmic/extracellular of THP-1 cells. Recommended concentration: 10µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB11513 (5 $\mu$ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB11513 Immunofluorescence analysis of paraformaldehyde fixed THP-1 cells immobilized on Shifix<sup>TM</sup> coverslip, permeabilized with 0.15% Triton. Primary incubation 1hr (10 $\mu$ g/ml) followed by Alexa Fluor 488 secondary antibody (2 $\mu$ g/ml), showing nuclear and cytoplasmic/extracellular staining. The nuclear stain is DAPI

(blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).