

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11513 - Goat Anti-S100A8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: 60B8AG, CAGA, calgranulin A, calgranulin-A, calprotectin L1L subunit, CFAG, CGLA, CP-10, cystic fibrosis antigen, L1Ag, leukocyte L1 complex light chain, MA387, MIF, migration inhibitory factor-related protein 8, MRP8, MRP-8, NIF, OTTHUMP00000015329, P8, protein S100-A8, S100 calcium binding protein A8, S100 calcium-binding protein A8 (calgranulin A), S100A8, urinary stone protein band A

Official Symbol: S100A8

Accession Number(s): NP_002955.2

Human GeneID(s): [6279](#)

Immunogen

Peptide with sequence C-KGADVWFKELDINT, from the internal region of the protein sequence according to NP_002955.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


IHC: Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nuclei and cytoplasmic/extracellular of THP-1 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB11513 (5 μ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB11513 Immunofluorescence analysis of paraformaldehyde fixed THP-1 cells immobilized on ShifixTM coverslip, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing nuclear and cytoplasmic/extracellular staining. The nuclear stain is DAPI

(blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).