



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06097 - Goat Anti-SAE1 / AOS1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SAE1, AOS1, SUA1, HSPC140, SUMO-1 activating enzyme subunit 1, sentrin/SUMO-activating protein AOS1, SUMO-1 activating enzyme E1 N subunit, ubiquitin-like protein SUMO-1 activating enzyme, SUMO1 activating enzyme subunit 1, FLJ3091, UBLE1A, activator of SUMO1, ubiquitin-like 1-activating enzyme E1A

Official Symbol: SAE1

Accession Number(s): NP_005491.1

Human GeneID(s): [10055](#)

Immunogen

Peptide with sequence MKGNGIVECLGPK, from the C Terminus of the protein sequence according to NP_005491.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx. 37kDa band observed in lysates of cell lines HEK293 and A549, and approx. 38-40kDa in lysates of cell line Jurkat (calculated MW of 38.4kDa according to NP_005491.1). Recommended concentration: 0.5-2µg/ml. Primary incubation 1 hour at room temperature.

Flow Cytometry: Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB06097 (1 μ g/ml) staining of HEK293 (A), A549 (B) and (2 μ g/ml) Jurkat (C) cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06097 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (1 μ g/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.