

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10681 - Goat Anti-SCOC Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SCOC, short coiled-coil protein, SCOCO, short coiled coil protein

Official Symbol: SCOC

Accession Number(s): NP_001146956.1; NP_001147135.1; NP_001147024.1;

NP_115936.2; NP_001147107.1; NP_001147162.1

Human GeneID(s): 60592

Non-Human GeneID(s): 56367 (mouse), 364981 (rat)

Important Comments: This antibody is expected to recognize all six reported isoforms

(NP_001146956.1; NP_001147135.1; NP_001147024.1; NP_115936.2;

NP_001147107.1; NP_001147162.1). Reported variants represent identical protein:

NP_001147057.1, NP_001146918.1, NP_115936.2

Immunogen

Peptide with sequence C-HTLEDLSARVDAVKE, from the internal region (near C Terminus) of the protein sequence according to NP_001146956.1; NP_001147135.1; NP_001147024.1; NP_115936.2; NP_001147107.1; NP_001147162.1.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 16kDa band observed in Human Skeletal Muscle lysates (calculated MW of 18.0kDa according to NP_001146956.1). Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10681 (0.1 μ g/ml) staining of Human Skeletal Muscle lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.