

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09387 - Goat Anti-SCRIB / Scribble Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SCRIB, scribbled homolog (Drosophila), CRIB1, SCRB1, SCRIB1, Vartul, scribble

**Official Symbol:** SCRIB

**Accession Number(s):** NP\_874365.3; NP\_056171.3

**Human GeneID(s):** [23513](#)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_874365.3; NP\_056171.3).

### Immunogen

Peptide with sequence C-PEGPGKEKELPGQ, from the internal region of the protein sequence according to NP\_874365.3; NP\_056171.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:1000.

**Western blot:** Approx 200-250kDa band observed in lysates of cell lines HeLa, HepG2 and U2OS (calculated MW of 178kDa according to NP\_874365.3). This molecular weight is routinely observed by other sources. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the cell junctions of MCF7 cells. Recommended concentration: 10µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB09387 (2 $\mu$ g/ml) staining of HeLa (A), HepG2 (B) and U2OS (C) cell lysate (35 $\mu$ g protein in RIPA buffer).  
Detected by chemiluminescence.



EB09387 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 $\mu$ g/ml) followed by Alexa Fluor 488 secondary antibody (2 $\mu$ g/ml), showing cell junction staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 $\mu$ g/ml) followed by Alexa

Fluor 488 secondary antibody (2ug/ml).