

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08933 - Goat Anti-SEPT7 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** SEPT7, septin 7, CDC10, CDC3, Nbla02942, SEPT7A, CDC10 (cell division cycle 10, S. cerevisiae, homolog), CDC10 cell division cycle 10 homolog, cell

division cycle 10

Official Symbol: SEPTIN7

Accession Number(s): NP\_001779.3; NP\_001011553.2; NP\_001229885.1

Human GeneID(s): 989

Non-Human GenelD(s): 235072 (mouse), 64551 (rat)

Important Comments: This antibody is expected to recognize all three reported isoforms

(NP\_001779.3; NP\_001011553.2; NP\_001229885.1).

# **Immunogen**

Peptide with sequence C-DNNKNKGQLTKSP, from the internal region of the protein sequence according to NP\_001779.3; NP\_001011553.2; NP\_001229885.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 45kDa band observed in nuclear lysates of cell line HeLa (calculated MW of 50.7kDa according to NP\_001779.3 and of 55.6kDa according to NP\_001011553.2). This molecular weight is routinely observed by other sources.

Recommended concentration: 0.3-1µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow, Horse



EB08933 (0.5 $\mu$ g/ml) staining of nuclear HeLa lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.