

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07013 - Goat Anti-SH2D1A/SLAM associated protein Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SH2D1A, SH2 domain protein 1A, Duncan\1s disease (lymphoproliferative syndrome), HGNC:10820, DSHP, EBVS, IMD5, LYP, MTCP1, SAP, XLP, XLPD, SH2 domain protein 1A, SLAM (signaling lymphocyte activation molecule) associated protein, T cell signal transduction molecule SAP, RP5-1052M9.3, FLJ18687, FLJ92177, Duncan's disease (lymphoproliferative syndrome), Duncan's disease, SLAM-associated protein, T cell signal transduction molecule SAP; signaling lymphocyte activation molecule-associated protein

Official Symbol: SH2D1A

Accession Number(s): NP_002342.1; NP_001108409.1

Human GeneID(s): 4068

Immunogen

Peptide with sequence C-QYPVEKKSSARSTQ, from the internal region of the protein sequence according to NP_002342.1; NP_001108409.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 16kDa band observed in Human Lymph Node lysates (calculated MW of 14.2kDa according to NP_002342.1).. Recommended concentration: 0.2-0.6µg/ml.

Immunoprecipitation:This antibody was deemed fit for IP under native conditions (observations from anonymous customer).

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Pig,



EB07013 (0.2 μ g/ml) staining of human lymph node lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.