

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

## EB05114 - Goat Anti-SHP2 / PTPN11 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** PTPN11, SHP2, protein tyrosine phosphatase, non-receptor type 11 (Noonan syndrome 1), CFC, NS1, BPTP3, PTP2C, SHP-2, PTP-1D, SH-PTP2, SH-PTP3, Noonan syndrome 1, protein-tyrosine phosphatase 2C, protein tyrosine phosphatase,

non-receptor type 11, MGC14433, protein tyrosine phosphatase-2

Official Symbol: PTPN11

Accession Number(s): NP\_002825.3; NP\_001317366.1

Human GeneID(s): 5781

Non-Human GenelD(s): 19247 (mouse), 25622 (rat)

### Immunogen

Peptide with sequence C-YENVGLMQQQKSFR, from the C Terminus of the protein sequence according to NP\_002825.3; NP\_001317366.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 70kDa band observed in Human Muscle lysates (calculated MW of 68.0kDa according to NP\_002825.3). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the nuclei and cytoplasm of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


# **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB05114 staining ( $2\mu g/ml$ ) of Human Muscle lysate (RIPA buffer, 35 $\mu g$  total protein per lane). Detected by chemiluminescence.



EB05114 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05114 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.