



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05292 - Goat Anti-SIAH1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SIAH1, seven in absentia homolog 1 (Drosophila), Siah-1, hSIAH1, HUMSIAH, Siah-1a, FLJ08065, seven in absentia homolog 1, siah E3 ubiquitin protein ligase 1, SIAH1A

Official Symbol: SIAH1

Accession Number(s): NP_003022.3; NP_001006611.1

Human GeneID(s): [6477](#)

Non-Human GeneID(s): 20437 (mouse), 140941 (rat)

Important Comments: This product is expected to recognize both reported isoforms (NP_003022.3; NP_001006611.1)

Immunogen

Peptide with sequence SRQTATALPTGTSC, from the N Terminus of the protein sequence according to NP_003022.3; NP_001006611.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx. 37kDa band observed in Human Liver lysates and approx. 30+37kDa bands in Rat Liver lysates (calculated MW of 34.6kDa according to Human NP_001006611.1 and 31.1kDa according to Rat NP_543181.2 and 38.2kDa according to Rat XP_038953364.1). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the nucleus and cytoplasm of U2OS and HepG2 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific References

This antibody (previous batch) has been successfully used in WB and IP on Human:

Liman J, Sroka K, Dohm CP, Deeg S, Bähr M, Kermer P.

Modulation of Huntingtin Toxicity by BAG1 is Dependent on an Intact BAG Domain.

Molecules 2010, 15, 6678-6687


PMID: 0

This antibody (previous batch) has been successfully used in Western blot on Drosophila:


Sroka K, Voigt A, Deeg S, Reed JC, Schulz JB, Bähr M, Kermer P.

BAG1 modulates huntingtin toxicity, aggregation, degradation, and subcellular distribution.


J Neurochem. 2009 Nov;111(3):801-7.
PMID: 19712056




EB05292 (1µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB05292 (1µg/ml) staining of RatLiver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB05292 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody.



EB05292 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody