

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10970 - Goat Anti-Sirtuin 3 (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Sirt3, sirtuin 3, sirtuin 3 (silent mating type information regulation 2, homolog) 3, 2310003L23Rik, Al848213, Sir2l3, NAD-dependent deacetylase sirtuin-3, OTTMUSP00000027441, OTTMUSP00000027444, OTTMUSP00000027445, SIR2-like

protein 3

Official Symbol: Sirt3

Accession Number(s): NP_071878.2; NP_001171275.1

Human GeneID(s): 23410

Non-Human GenelD(s): 64384 (mouse), 293615 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_071878.2; NP_001171275.1). Reported variants represent identical proteins: NP_071878.2; NP_001120823.1

Immunogen

Peptide with sequence C-RASGIPASKLVEAH, from the internal region of the protein sequence according to NP_071878.2; NP_001171275.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


Western blot: Approx 30kDa band observed in Mouse Heart lysates (calculated MW of 28.8kDa according to NP_071878.2). An additional band of unknown identity was also consistently observed at 15kDa. This band was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature. Preliminary testing was unsuccessful on Rat Kidney for this particular batch.

Flow Cytometry: Flow cytometric analysis of NIH3T3 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Mouse


Expected from sequence similarity: Human, Mouse, Rat, Pig



EB10970 (0.3 μ g/ml) staining of Mouse Heart lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB10970 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB10970 Flow cytometric analysis of paraformaldehyde fixed NIH3T3 cells (blue line), permeabilized with 0.5% Triton, showing a low level of staining. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.