

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08818 - Goat Anti-SLC10A2 / ASBT Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SLC10A2, ASBT, solute carrier family 10 (sodium/bile acid cotransporter family), member 2, ISBT, NTCP2, OTTHUMP00000040691, ileal apical sodium-dependent bile acid transporter, ileal sodium-dependent bile acid transporter

**Official Symbol:** SLC10A2

**Accession Number(s):** NP\_000443.1

**Human GeneID(s):** [6555](#)

### Immunogen

Peptide with sequence C-YKANGGFQPDEK, from the C Terminus of the protein sequence according to NP\_000443.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.


**Western blot:** Preliminary experiments gave an approx. 60kDa band in Human Kidney lysates after 0.1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 37.7kDa according to NP\_000443.1. The 60kDa band was successfully blocked by incubation with the immunizing peptide.

**IHC:** In paraffin embedded Human Small Intestine shows staining of cells in the lamina propria between the intestinal villi. Recommended concentration: 3-6µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB08818 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.