

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05131 - Goat Anti-SLC16A7 / MCT2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SLC16A7, solute carrier family 16 (monocarboxylic acid transporters), member 7, MCT2, monocarboxylate transporter 2, solute carrier family 16, member 7 (monocarboxylic acid transporter 2), solute carrier family 16, member 7

**Official Symbol:** SLC16A7

**Accession Number(s):** NP\_004722.2

**Human GeneID(s):** [9194](#)

### Immunogen

Peptide with sequence C-SNAQSVTSERETNI, from the C Terminus of the protein sequence according to NP\_004722.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.


**Western blot:** Approx 40-45kDa band observed in A549, lung epithelial (lung carcinoma) lysates (calculated MW of 52.2 kDa according to NP\_004722). This molecular weight is routinely observed by other sources. Recommended concentration: 1-3µg/ml.

**IHC:** Paraffin embedded Human Testis. Recommended concentration: 2.5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB05131 (1 $\mu$ g/ml) staining of A549 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB05131 (2.5 $\mu$ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.