

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05131 - Goat Anti-SLC16A7 / MCT2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SLC16A7, solute carrier family 16 (monocarboxylic acid transporters), member 7, MCT2, monocarboxylate transporter 2, solute carrier family 16, member 7

(monocarboxylic acid transporter 2), solute carrier family 16, member 7

Official Symbol: SLC16A7

Accession Number(s): NP_004722.2

Human GeneID(s): 9194

Immunogen

Peptide with sequence C-SNAQSVTSERETNI, from the C Terminus of the protein sequence according to NP_004722.2.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 40-45kDa band observed in A549, lung epithelial (lung carcinoma) lysates (calculated MW of 52.2 kDa according to NP_004722). This molecular weight is

routinely observed by other sources. Recommended concentration: 1-3μg/ml. **IHC:** Paraffin embedded Human Testis. Recommended concentration: 2.5μg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB05131 (1 μ g/ml) staining of A549 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB05131 (2.5 μ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.