

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11236 - Goat Anti-Slc5a1 (rat) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: high affinity sodium-glucose cotransporter, MGC93553, Na(+)/glucose cotransporter 1, Sglt1, sodium/glucose cotransporter 1, solute carrier family 5 (sodium/glucose cotransporter), member 1, solute carrier family 5, member 1, Solute carrier family 5, member alpha 1 (Na+/glucose cotransporter), Slc5a1

Official Symbol: Slc5a1

Accession Number(s): NP_037165.1 Non-Human GeneID(s): 25552 (rat)

Immunogen

Peptide with sequence C-KDTIEIDAEAPQKEK, from the internal region of the protein sequence according to NP_037165.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 70kDa band observed in Rat Kidney lysates (calculated MW of 72.1kDa according to NP_037165.1). Recommended concentration: 0.3-2μg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Rat Kidney. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Rat


Expected from sequence similarity: Rat

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB11236 (1μg/ml) staining of Rat Kidney lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB11236 (10μg/ml): Staining of paraffin embedded Rat Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-Staining (1:500).



 ${\tt EB11236\ Negative\ Control\ showing\ staining\ of\ paraffin\ embedded\ Rat\ Kidney,\ with\ no\ primary\ antibody.}$