



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06217-B - Goat Anti-SNX5, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SNX5, sorting nexin 5

**Official Symbol:** SNX5

**Accession Number(s):** NP\_055241.1; NP\_001269383.1

**Human GeneID(s):** [27131](#)

**Non-Human GeneID(s):** 69178 (mouse), 296199 (rat)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_055241.1; NP\_001269383.1). Reported variants represent identical protein: NP\_055241.1, NP\_689413.1

### Immunogen

Peptide with sequence SLLQSCIDLFKNN., from the C Terminus of the protein sequence according to NP\_055241.1; NP\_001269383.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 48kDa band observed in lysates of cell line A549 (calculated MW of 46.8kDa according to NP\_055241.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow



Biotinylated EB06217 (2µg/ml) staining of A549 lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.