

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11971 - Goat Anti-Somatostatin proprotein (aa58-70) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SST, somatostatin, SMST, growth hormone release-inhibiting factor,

somatostatin-14, somatostatin-28

Official Symbol: SST

Accession Number(s): NP_001039.1

Human GeneID(s): 6750

Non-Human GenelD(s): 20604 (mouse), 24797 (rat)

Immunogen

Peptide with sequence C-SEPNQTENDALEP, from the internal region of the protein sequence according to NP_001039.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:1000.

Immunofluorescence/ Immunocytochemistry: Strong expression of the protein seen in the plasma membrane and endoplasmic reticulum of U2OS and U251 cells.

Recommended concentration: $10\mu g/ml$. This antibody has been used successfully in ICC on Mouse.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

Specific Reference


This antibody has been successfully used in ICC on Mouse:

Anna Billeschou Bomholt, Christian Dall Johansen, Jens Bager Christensen, Sasha Alexandra Sampson Kjeldsen, Katrine Douglas Galsgaard, Marie Winther-Sørensen, Reza Serizawa, Mads Hornum, Esteban Porrini, Jens Pedersen, Cathrine Ørskov, Lise Lotte Gluud, Charlotte Mehlin Sørensen, Jens Juul Holst, Reidar Albrechtsen & Nicolai Jacob Wewer Albrechtsen


Evaluation of commercially available glucagon receptor antibodies and glucagon receptor expression.

Commun Biol. 2022 Nov 22;5(1):1278.

PMID: 36418521



EB11971 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing plasma membrane and endoplasmic reticulum staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB11971 Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing plasma membrane and endoplasmic reticulum staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).