



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07378 - Goat Anti-SOX2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SOX2, SRY (sex determining region Y)-box 2, ANOP3, MGC2413, SRY-related HMG-box gene 2, sex-determining region Y-box 2, transcription factor SOX2

Official Symbol: SOX2

Accession Number(s): NP_003097.1

Human GeneID(s): [6657](#)

Immunogen

Peptide with sequence YNMMETELKPPGPQQ-C, from the N Terminus of the protein sequence according to NP_003097.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 40kDa band observed in HEK293 and MCF7 cell lysates and in nuclear lysates of cell line MCF7 (calculated MW of 34.3kDa according to JNP_003097.1).

This molecular weight is routinely observed by other sources. Recommended concentration: 0.1-0.3µg/ml. Primary incubation 1 hour at room temperature.


IHC: Paraffin embedded Human Tonsil. Recommended concentration: 6-8µg/ml.

Flow Cytometry: Flow cytometric analysis of HEK293 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow




EB07378 (0.3 μ g/ml) staining of MCF7 (A) HEK293 (B) and MCF7 nuclear (C) cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB07378 Flow cytometric analysis of paraformaldehyde fixed HEK293 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (1 μ g/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB07378 (8µg/ml) staining of paraffin embedded Human Tonsil. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07378 Negative Control showing staining of paraffin embedded Human Tonsil, with no primary antibody.