



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB13088 - Goat Anti-SARS-COV2 Spike protein-S1 Domain (N Terminal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) isolate Wuhan-Hu-1, Spike glycoprotein, E2, Peplomer protein

Official Symbol: SARS-CoV-2

Accession Number(s): P0DTC2; NC_045512.2/MN908947

Immunogen

Peptide with sequence PROPRIETARY-SARS-CoV-2, from the N Terminus of the protein sequence according to P0DTC2; NC_045512.2/MN908947.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


Western blot: Approx 100-110kDa band observed on recombinant SARS-COV2 Spike Protein S1. Recommended concentration 0.5-1ug/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Transient expression of SARS-CoV-2 spike protein expressing plasmid, seen in the cytoplasm of HEK293 transfected cells.


Species Reactivity

Tested: Viral

Expected from sequence similarity: Human Coronavirus



EB13088 (1 μ g/ml) staining of recombinant Spike Protein S1 (1 μ g protein in PBS, pH 7.4). Detected by chemiluminescence.



EB13088 Immunofluorescence analysis of paraformaldehyde fixed HEK293 cells transiently transfected with SARS-CoV-2 spike protein expressing plasmid, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: non-transfected HEK293 cells.