



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05341 - Goat Anti-Spinesin / TMPRSS5 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TMPRSS5, SPINESIN, transmembrane protease, serine 5 (spinesin), MGC141886, MGC148044, SPINESIN

Official Symbol: TMPRSS5

Accession Number(s): NP_110397.2

Human GeneID(s): [80975](#)

Immunogen

Peptide with sequence C-LDWIHDTAQDSLL, from the C Terminus of the protein sequence according to NP_110397.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Experiments gave bands at approx 50kDa and 29kDa in Human Brain (Cerebral Cortex). These bands correspond to earlier findings in literature with different antibodies (Yamaguchi et al, J Biol Chem. 2002 Mar 1;277(9):6806-12; PMID: 11741986). This protein has a calculated MW of 49.6kDa according to NP_110397.1. Recommended concentration: 0.5-1.5µg/ml.

IHC: Paraffin embedded Human Liver and Small Intestine. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Dog



EB05341 staining (0.5µg/ml) of Human Brain (Cerebral Cortex) lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



EB05341 (5µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05341 (5µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.