

#### **International Office**

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06432 - Goat Anti-Sprouty Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: SPRY1, hSPRY1, sprouty homolog 1, antagonist of FGF signaling (Drosophila), sprouty (Drosophila) homolog 1 (antagonist of FGF signaling), sprouty, Drosophila, homolog of, 1 (antagonist of FGF signaling) Official Symbol: SPRY1 Accession Number(s): NP\_005832.1; NP\_955359.1 Human GenelD(s): <u>10252</u> Important Comments: NP\_005832.1 and NP\_955359.1 represent varients of the same protein.

#### Immunogen

Peptide with sequence CPSRGQGKPS, from the C Terminus of the protein sequence according to NP\_005832.1; NP\_955359.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 37kDa band observed in Human Kidney lysates (calculated MW of 35.1kDa according to NP\_005832.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature. This.product has been successfully used in ERMS cells (PMID: 20068162). Recommended concentration: 1-3µg/ml.

**IHC:** This product has been successfully used by customers in paraffin embedded Mouse Brain, showing nuclear staining. Recommended concentration: 1-3µg/ml.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm and Golgi apparatus of HeLa and HepG2 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HEK293 cells. Recommended concentration: 10ug/ml.

# **Species Reactivity**

Tested: Human, Mouse Expected from sequence similarity: Human



EB06432 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic and Golgi apparatus staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).





EB06432 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic and Golgi apparatus staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB06432 Flow cytometric analysis of paraformaldehyde fixed HEK293 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (0.4ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.