

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05127 - Goat Anti-STAP2 / BKS Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: STAP2, BKS, STAP-2, FLJ20234, breast tumor kinase substrate, brk kinase substrate, signal-transducing adaptor protein-2, signal transducing adaptor family member 2

Official Symbol: STAP2

Accession Number(s): NP_060190.2; NP_001013863.1

Human GeneID(s): [55620](#)

Important Comments: This antibody is expected to recognise isoforms 1 (NP_060190.2) and isoform 2 (NP_001013863.1).

Immunogen

Peptide with sequence C-ELQKKLEKRRALEH, from the C Terminus of the protein sequence according to NP_060190.2; NP_001013863.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 48kDa band observed in Human Heart lysates and 45kDa band observed in Human Breast Lysates (calculated MW of 49.6kDa according to NP_060190 and of NP_001013863.1). Recommended concentration: 0.5-1.5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

Specific References

This antibody (previous batch) has been successfully used in Western blot on Human:

Ikeda O, Mizushima A, Sekine Y, Yamamoto C, Muromoto R, Nanbo A, Oritani K, Yoshimura A, Matsuda T.

Involvement of STAP-2 in Brk-mediated phosphorylation and activation of STAT5 in breast cancer cells.

Cancer Sci. 2011 Apr;102(4):756-61.

PMID: 21205088

This antibody (previous batch) has been successfully used in Western blot on Human:

Ikeda O, Sekine Y, Mizushima A, Nakasuji M, Miyasaka Y, Yamamoto C, Muromoto R, Nanbo A, Oritani K, Yoshimura A, Matsuda T.

Interactions of STAP-2 with Brk and STAT3 participate in cell growth of human breast cancer cells.

J Biol Chem. 2010 Dec 3;285(49):38093-103.

PMID: 20929863

This antibody (previous batch) has been successfully used in WB and IF on Human:

Sekine Y, Ikeda O, Tsuji S, Yamamoto C, Muromoto R, Nanbo A, Oritani K, Yoshimura A, Matsuda T.

Signal-transducing adaptor protein-2 regulates stromal cell-derived factor-1 alpha-induced chemotaxis in T cells.

J Immunol. 2009 Dec 15;183(12):7966-74.

PMID: 19933863


This antibody (previous batch) has been successfully used in the following paper:

Ikeda O, Miyasaka Y, Sekine Y, Mizushima A, Muromoto R, Nanbo A, Yoshimura A, Matsuda T.

STAP-2 is phosphorylated at tyrosine-250 by Brk and modulates Brk-mediated STAT3 activation.

Biochem Biophys Res Commun. 2009 Jun 19;384(1):71-5.

PMID: 19393627



EB05127 (0.5µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.