

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07108 - Goat Anti-Stearoyl-CoA desaturase Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SCD, SCD1, HGNC:10571, FADS5, PRO0998, acyl-CoA desaturase, delta-9-desaturase, fatty acid desaturase, predicted protein of HQ0998, stearoyl-CoA desaturase

Official Symbol: SCD

Accession Number(s): NP_005054.3

Human GeneID(s): [6319](#)

Immunogen

Peptide with sequence C-RIKRTGDGNYKSG, from the C Terminus of the protein sequence according to NP_005054.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Preliminary experiments gave an approx 30kDa band in Human Adipose, Brain and Liver lysates after 0.3 µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 41.5kDa according to NP_005054.3. The 30kDa band was successfully blocked by incubation with the immunizing peptide.


IHC: Paraffin embedded Human Liver and Colon. Recommended concentration: 5µg/ml.

Flow Cytometry: Exclusively staining of HepG2 when mixed with Peripheral Blood Lymphocytes.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human



EB07108 (5µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07108 (5µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07108 (2.5ul) staining of HepG2 cells mixed in with PBL. Detected by fluorescence with PE for SCD in green and with OC515 for CD45 (CYT-450C) in red.