



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09416 - Goat Anti-STK39 / SPAK Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: STK39, serine threonine kinase 39 (STE20/SPS1 homolog, yeast), DCHT, DKFZp686K05124, PASK, SPAK, Ste20-like protein kinase, proline-alanine-rich STE20-related kinase, small intestine SPAK-like kinase

Official Symbol: STK39

Accession Number(s): NP_037365.2

Human GenelD(s): [27347](#)

Non-Human GenelD(s): 53416 (mouse), 54348 (rat)

Immunogen

Peptide with sequence C-SQEKSRRVKEENPE, from the internal region of the protein sequence according to NP_037365.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.


Western blot: Approx 65kDa band observed in Human Brain (Cerebellum) lysate (calculated MW of 59.5kDa according to NP_037365.2). The observed molecular weight corresponds to earlier findings with different antibodies from other commercial sources. Recommended concentration: 0.3-1µg/ml.

IHC: Paraffin embedded Human Testis. Recommended concentration: 6µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog



EB09416 (0.3 μ g/ml) staining of Human Cerebellum lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB09416 (6 μ g/ml) staining of paraffin embedded Human Testis. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB09416 Negative Control showing staining of paraffin embedded Human Testis, with no primary antibody.