

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06552 - Goat Anti-Swiprosin 1 / EFHD2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EFHD2, MGC4342, EF hand domain containing 2

Official Symbol: EFHD2

Accession Number(s): NP_077305.2

Human GeneID(s): 79180

Non-Human GenelD(s): 27984 (mouse)

Important Comments: This antibody does not cross-react with Mouse Swiprosin 2. Data

kindly provided by Sebastian Duetting and Dirk Mielenz.

Immunogen

Peptide with sequence CAAFKELQSTFK, from the C Terminus of the protein sequence according to NP_077305.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 30kDa band observed in Human Thymus, Human Lymph node and

Human Tonsil lysates (calculated MW of 26.7kDa according to NP_077305.2). Recommended concentration: 0.5-1.5 μ g/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow


Specific Reference

This antibody (previous batch) has been successfully used in Western blot on Human:

Brachs S, Lang C, Buslei R, Purohit P, Fürnrohr B, Kalbacher H, Jäck HM, Mielenz D. Monoclonal antibodies to discriminate the EF hand containing calcium binding adaptor proteins EFhd1 and EFhd2.

Monoclon Antib Immunodiagn Immunother. 2013 Aug;32(4):237-45.

PMID: 23909416



EB06552 (0.5μg/ml) staining of Human Thymus, Human Lymph node and Human Tonsil lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.