

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09721 - Goat Anti-TAF8 / TAFII43 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** TAF8, TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 43kDa, 43, FLJ32821, II, TAF, TAFII43, TBN, OTTHUMP00000016392, TAF(II)43, TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 50 kD, TATA box binding protein (TBP)-associated factor, RNA polymerase II, A, 45/50kDa, TBP-associated factor 8, TBP-associated factor TAFII43, TBP-associated factor, RNA polymerase II, 43 kD, taube nuss homolog

**Official Symbol:** TAF8

**Accession Number(s):** NP\_612639.2

**Human GeneID(s):** [129685](#)

**Non-Human GeneID(s):** 63856 (mouse), 316216 (rat)

### Immunogen

Peptide with sequence KSYCEHTARTQPT, from the internal region of the protein sequence according to NP\_612639.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.


**Western blot:** Approx. 37kDa band observed in Human Hippocampus lysates, approx. 38kDa in Mouse and Rat Brain lysates and approx. 35kDa in lysates of cell line NIH3T3 (calculated MW of 34.3kDa according to Human NP\_612639.2, 34.0kDa according to Mouse NP\_071298.1 and 34.1kDa according to Rat NP\_001101667.1). Recommended concentration: 0.5-µg/ml. Primary incubation 1 hour at room temperature.

**Flow Cytometry:** Flow cytometric analysis of NIH3T3 cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human, Mouse, Rat


**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB09721 (0.5µg/ml) staining of NIH3T3 cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB09721 (1µg/ml) staining of Human Hippocampus (A), Rat Brain (B) and (2ug/ml) Mouse Brain (C) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB09721 Flow cytometric analysis of paraformaldehyde fixed NIH3T3 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.