



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06281 - Goat Anti-SM22 Alpha / Transgelin Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TAGLN, SM22, SMCC, WS3-10, transgelin, SM22-alpha, DKFZp686P11128, TAGLN1, DKFZp686B01212, transgelin variant 2

Official Symbol: TAGLN

Accession Number(s): NP_003177.2

Human GeneID(s): [6876](#)

Non-Human GeneID(s): 21345 (mouse), 25123 (rat)

Important Comments: NP_003177.2 and NP_001001522.1 represent identical protein.

The transgelin protein is found in fibroblasts and smooth muscle and is known as a transformation or shape change protein that is involved with actin cross-linking. Transgelin is one of the earliest markers of differentiated smooth muscle cells and recent evidence suggests that transgelin acts as a tumour suppressor.

Immunogen

Peptide with sequence C-MTGYGRPRQIIS, from the C Terminus of the protein sequence according to NP_003177.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 24kDa band observed in Human Skeletal Muscle and Testis lysates, and approx 25kDa in Mouse Skeletal Muscle lysates (calculated MW of 22.6kDa according to Human NP_003177.2 and Mouse NP_035656.1). These molecular weights are routinely observed by other sources. Recommended concentration: 0.003-0.03 µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific References

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Angela Corvino et al.

Antagonizing S1P3 Receptor with Cell-Penetrating Peptides in Skeletal Muscle Fibrosis
Int J Mol Sci. 2021 Aug; 22(16): 8861.
PMID: 34445567

This antibody (previous batch) has been successfully used in Western blot on Human:

Caterina Bernacchioni, Tommaso Capezzuoli, Valentina Vannuzzi, Francesca Malentacchi, Francesca Castiglione, Francesca Cencetti, Marcello Ceccaroni, Chiara

Donati, Paola Bruni, Felice Petraglia

Sphingosine 1-phosphate receptors are dysregulated in endometriosis: possible implication in transforming growth factor β -induced fibrosis

Fertil Steril. 2020 Sep 6;S0015-0282(20)30766-4.

PMID: 32907751

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Bruno G, Cencetti F, Pertici I, Japtok L, Bernacchioni C, Donati C, Bruni P.

CTGF/CCN2 exerts profibrotic action in myoblasts via the up-regulation of sphingosine kinase-1/S1P3 signaling axis: Implications in the action mechanism of TGF β .

Biochim Biophys Acta. 2015 Feb;1851(2):194-202.

PMID: 25457224

This antibody (previous batch) has been successfully used in Western blot on Human:

Donati C, Marseglia G, Magi A, Serrati S, Cencetti F, Bernacchioni C, Nannetti G, Benelli M, Brunelli S, Torricelli F, Cossu G, Bruni P.

Sphingosine 1-phosphate induces differentiation of mesoangioblasts towards smooth muscle. A role for GATA6.

PLoS One. 2011;6(5):e20389.

PMID: 21629665

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Cencetti F, Bernacchioni C, Nincheri P, Donati C, Bruni P.

Transforming growth factor-beta1 induces transdifferentiation of myoblasts into myofibroblasts via up-regulation of sphingosine kinase-1/S1P3 axis.

Mol Biol Cell. 2010 Mar 15;21(6):1111-24.

PMID: 20089836

This antibody (previous batch) has been successfully used in Western blot on Human:

Nincheri P, Luciani P, Squecco R, Donati C, Bernacchioni C, Borgognoni L, Luciani G, Benvenuti S, Francini F, Bruni P.


Sphingosine 1-phosphate induces differentiation of adipose tissue-derived mesenchymal stem cells towards smooth muscle cells.

Cell Mol Life Sci. 2009 May;66(10):1741-54.

PMID: 19337690



EB06281 (0.03 μ g/ml) staining of Human Skeletal muscle (A) and (0.003 μ g/ml) Testes (B) lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06281 (0.03 μ g/ml) staining of Mouse Skeletal muscle lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.