



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB02504 - Goat Anti-TANK2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TNKS2, tankyrase, TRF1-interacting ankyrin-related ADP-ribose polymerase 2, PARP-5b, PARP-5c, PARP5B, PARP5C, TANK2, TNKL, tankyrase 2

Official Symbol: TNKS2

Accession Number(s): NP_079511.1

Human GeneID(s): [80351](#)

Important Comments: Fusion protein containing 73 aas from region aa800-900 (according to NP_079511.1). No cross reactivity expected with TANK1.

Immunogen

Fusion protein containing 73 aas from aa800-900 region of the protein sequence according to NP_079511.1. No crossreactivity expected with TANK1.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


Applications Tested

Western blot: In transfected HEK293 transiently expressing TANK2 a band of approx. 127kDa is observed. This band is not observed in the non-transfected HEK293. The calculated molecular size is 127 kDa according to NP_079511.1. Recommended concentration, 0.5-1.5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



HEK293 overexpressing TANK2 (lane 4) and TANK1 (lane 2) and probed with EB02504 (mock transfection in first lane). Lane three is empty. Lower panel shows the same lysates probed for alpha-Actin to show protein levels. Primary incubation (0.5ug/ml) was overnight at 4°C. Detected by chemiluminescence.