

### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08220 - Goat Anti-Tankyrase 2 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** TNKS2, tankyrase, TRF1-interacting ankyrin-related ADP-ribose polymerase 2, PARP-5b, PARP-5c, PARP5B, PARP5C, TANK2, TNKL, tankyrase 2

Official Symbol: TNKS2

Accession Number(s): NP\_079511.1

Human GeneID(s): 80351

Non-Human GenelD(s): 74493 (mouse)

### **Immunogen**

Peptide with sequence C-HRRKEVSEENHNH, from the internal region of the protein sequence according to NP\_079511.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx. 125 kDa band observed in lysates of cell line Jurkat (calculated MW of 127 kDa according to NP\_079511.1). Recommended concentration: 0.5-1.5μg/ml. An additional band of unknown identity was also consistently observed at 45 kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Dog



EB08220 (0.5 $\mu$ g/ml) staining of Jurkat lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.