

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06265 - Goat Anti-TIGD1 / EEYORE Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** TIGD1, EEYORE, tigger transposable element derived 1, jerky (mouse) homolog-like, hypothetical protein LOC200765, jerky homolog-like

Official Symbol: TIGD1

Accession Number(s): NP\_663748.1

Human GenelD(s): 200765

#### **Immunogen**

Peptide with sequence C-PAKRVRLTEGSD, from the C Terminus of the protein sequence according to NP\_663748.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Preliminary experiments gave no signal but low background in Human

Brain and Kidney extracts at up to 1µg/ml.


Immunofluorescence: Strong expression of the protein seen in the nuclei of U2OS cells.

Recommended concentration: 10µg/ml.

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB06265 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).