

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# **EB05048 - Goat Anti-Tissue Factor Pathway Inhibitor Antibody**

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** tissue factor pathway inhibitor (lipoprotein-associated coagulation inhibitor), TFPI, EPI, LACI, TFI, TFPI1, OTTHUMP00000163472,

OTTHUMP00000205431, anti-convertin, extrinsic pathway inhibitor, lipoprotein-associated coagulation inhibitor, tissue factor pathway inhibitor

Official Symbol: TFPI

Accession Number(s): NP\_006278.1

Human GenelD(s): 7035

Important Comments: This antibody is expected to recognize isoform a (NP\_006278.1)

only.

### **Immunogen**

Peptide with sequence C-VKIAYEEIFVKNM, from the C Terminus of the protein sequence according to NP\_006278.1.

Please note the peptide is available for sale.

#### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 50kDa band observed in lysates of cell lines HepG2 and HeLa (calculated MW of 35.0kDa according to NP\_006278.1). In transfected HEK293 transiently expressing TFPI bands of approx. 38kDa to 45kDa are observed. These bands are not observed in the non-transfected HEK293. Recommended concentration: 0.3-1µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



HEK293 overexpressing TFPI (RC219033) and probed with EB05048 (mock transfection in first lane), tested by Origene.



EB05048 staining (0.3μg/ml) of Hela lysate (RIPA buffer, 35μg total protein cells per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.