

International Office

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06556 - Goat Anti-TOM1L2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TOM1L2, FLJ32746, target of myb1-like 2 (chicken), target of myb1 (chicken) homolog-like 1, target of myb1 (chicken) homolog-like 2 Official Symbol: TOM1L2 Accession Number(s): NP_001028723.1; NP_001076437.1 Human GeneID(s): <u>146691</u> Important Comments: This antibody is expected to recognise both reported isoforms.

Immunogen

Peptide with sequence CRKKPERSEDALFAL, from the C Terminus of the protein sequence according to NP_001028723.1; NP_001076437.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 60kDa band observed in Human Fibroblast A431 lysates (calculated MW of 51.8kDa according to NP_653279). Recommended concentration: 0.1-1µg/ml.

IHC: In paraffin embedded Human Skeletal Muscle shows variable staining from fibre to fibre. Recommended concentration, 5-10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Dog

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa 20kDa
15kDa

EB06556 (0.03µg/ml) staining of A431 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB06556 (5µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.